

PAN056Mu01

Polyclonal Antibody

To T-Cell Immunoreceptor With Ig And ITIM Domains Protein (TIGIT)

Organism Species: Mus musculus (Mouse)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

[PRODUCT INFORMATION]

Immunogen: TIGIT, Mouse

Clonality: Polyclonal

Host: Rabbit


Immunoglobulin Type: IgG

Purification: Affinity Chromatography.

Applications: WB, ICC, IHC-P, IHC-F, ELISA

Concentration: 200µg/mL

UOM: 100µg



Sample: Recombinant TIGIT, Mouse

[IMMUNOGEN INFORMATION]

Immunogen: Recombinant TIGIT (Gly28~Ser246) expressed in *E.coli*.

Accession No.: RPN056Mu01

Sequence: The target protein is fused with two N-terminal Tags, His-tag and T7-tag and its sequence is listed below.

MGSSHHHHHHH SSGLVPRGSH MASMTGGQQM GRGSEF-GTI DTKRNISAE
GGSVILQCHF SSDTA EVTQV DWKQQDQLLA IYSVDLGWHV ASVFSDRVVP GPSLGLTFQS
LTMNDTGEYF CTYHTYPGGI YKGRIFLKVQ ESSDDRNGLA QFQTAPLGGT MAAVLGLICL
MVTGVTVLAR KDKSIRMHSI ESGLGRTEAE PQEWNLRSLSPGSPVQTQT APAGPCGEQA
EDDYADPQEY FNVLSYRSLE SFI AVS

[ANTIBODY SPECIFICITY]

The antibody is a rabbit polyclonal antibody raised against TIGIT. It has been selected for its ability to recognize TIGIT in immunohistochemical staining and western blotting.

[APPLICATIONS]

Western blotting: 1:50-400

Immunocytochemistry in formalin fixed cells: 1:50-500

Immunohistochemistry in formalin fixed frozen section: 1:50-500

Immunohistochemistry in paraffin section: 1:10-100

Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

[CONTENTS]

Form & Buffer: Supplied as solution form in PBS, pH7.4, containing 0.02% NaN₃, 50% glycerol.

[QUALITY CONTROL]

Content: The quality control contains recombinant TIGIT (Gly28~Ser246) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.
5uL per well when used in enhanced chemiluminescent (ECL).

Note: The quality control is specifically manufactured as the positive control. Not used for other purposes.

Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN₃ 0.02%.

[STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.