

PAA101Ra02

Polyclonal Antibody to Matrix Metalloproteinase 3 (MMP3)

Organism Species: Rattus norvegicus (Rat)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

[PRODUCT INFORMATION]

Immunogen: MMP3, Rat **Purification:** Affinity Chromatography.

Clonality: Polyclonal Applications: WB, ICC, IHC-P, IHC-F, ELISA

Host: Rabbit Concentration: 200µg/mL

Immunoglobulin Type: IgG **UOM**: 100μg

[IMMUNOGEN INFORMATION]

Immunogen: Recombinant MMP3 (Phe98~Cys475) expressed in *E.coli*.

Accession No.: RPA101Ra02

Sequence: The target protein is fused with two N-terminal Tags, His-tag and

T7-tag and its sequence is listed below.

MGSSHHHHHH SSGLVPRGSH MASMTGGQQM GRGSEF- FST FPGSPKWRKN HISYRIVNYT LDLPRESVDS AIERALKVWE EVTPLTFSRI SEGEADIMIS FAVEEHGDFI PFDGPGMVLA HAYAPGPGTN GDAHFDDDER WTDDVTGTNL FLVAAHELGH SLGLFHSANA EALMYPVYKS STDLARFHLS QDDVDGIQSL YGPPTESPDV LVVPTKSNSL DPETLPMCSS ALSFDAVSTL RGEVLFFKDR HFWRKSLRTP EPGFYLISSF WPSLPSNMDA AYEVTNRDTV FILKGNQIWA IRGHEELAGY PKSIHTLGLP ETVQKIDAAI SLKDQKKTYF FVEDKFWRFD EKKQSMDPEF PRKIAENFPG IGTKVDAVFE AFGFLYFFSG SSQLEFDPNA GKVTHILKSN SWFNC

[ANTIBODY SPECIFITY]

The antibody is a rabbit polyclonal antibody raised against MMP3. It has been selected for its ability to recognize MMP3 in immunohistochemical staining and western blotting.

[APPLICATIONS]

Western blotting: 1:100-400


Immunocytochemistry in formalin fixed cells: 1:100-500

Immunohistochemistry in formalin fixed frozen section: 1:100-500


Immunohistochemistry in paraffin section: 1:50-200 Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

[IMAGES]



Used in DAB staining on fromalin fixed paraffin-embedded Kidney tissue



Used in Western Blot, Sample:
Recombinant MMP3, Rat

[CONTENTS]

Form & Buffer: Supplied as solution form in PBS, pH7.4, containing 0.02% NaN₃, 50% glycerol.

[QUALITY CONTROL]

Content: The quality control contains recombinant MMP3 (Phe98~Cys475) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemilumescent (ECL).

Note: The quality control is specifically manufactured as the positive control. Not used for other purposes.

Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN₃ 0.02%.

[STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.