

PAA326Ra01

Polyclonal Antibody to Sp100 Nuclear Antigen (Sp100)
Organism Species: Rattus norvegicus (Rat)

Instruction manual

←28kDa

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

Western Blot

44

33

26

10

[PRODUCT INFORMATION]

Immunogen: Sp100, Rat

Clonality: Polyclonal

Host: Rabbit

Immunoglobulin Type: IgG

Purification: Affinity Chromatography.

Applications: WB, ICC, IHC-P, IHC-F, ELISA

Concentration: 200µg/mL

UOM: 100µg

22 18 ography.

Sample: Recombinant Sp100, Rat

[IMMUNOGEN INFORMATION]

Immunogen: Recombinant Sp100 (Phe116~Pro333) expressed in E.coli.

Accession No.: RPA326Ra01

Sequence: The target protein is fused with two N-terminal Tags, His-tag and

T7-tag and its sequence is listed below.

MGSSHHHHHH SSGLVPRGSH MASMTGGQQM GRGSEF- FERVL PNGLCFQRSD GGDPNSQLSL EQGPNGSCSQ ATLTLSPFNH SFSDGWGSND RLNSSLTHGN QSENHQLPGS PNHLGAVQLN HDLQINPCSV RLVDIKKENI SYSLDDNQQT QARTNENQGS EIIELSSDED SNDEINLSEA STLILSKPET AHSRQPPTPR TYRRGDTSDT SSSSLDRRRK RTGRMLSSSS SELSSGEESE EFLRSGSGRL GCP



[ANTIBODY SPECIFITY]

The antibody is a rabbit polyclonal antibody raised against Sp100. It has been selected for its ability to recognize Sp100 in immunohistochemical staining and western blotting.

[APPLICATIONS]

Western blotting: 1:100-400

Immunocytochemistry in formalin fixed cells: 1:100-500

Immunohistochemistry in formalin fixed frozen section: 1:100-500

Immunohistochemistry in paraffin section: 1:50-200 Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

[CONTENTS]

Form & Buffer: Supplied as solution form in PBS, pH7.4, containing 0.02% NaN₃, 50% glycerol.

[QUALITY CONTROL]

Content: The quality control contains recombinant Sp100 (Phe116~Pro333) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemilumescent (ECL).

Note: The quality control is specifically manufactured as the positive control. Not used for other purposes.

Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN₃ 0.02%.

[STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.